

JASON'S QUESTIONS AND SOLUTIONS

QUESTION 1: Solve $x^2 - 4 \leq 0$.

SOLUTION

STEPS


1. Get all terms on one side and zero on the other side.
2. Solve the corresponding equation to get the roots α , β .
3. Carry out the region test. Use the roots in ascending order to form regions: $\leftarrow \alpha \leftrightarrow \beta \rightarrow$ Choose a nice number in each region to test the inequality using the **test box**.
4. Based on the region test write down the solutions.

1. $x^2 - 4 \leq 0$

2. Solve $x^2 - 4 = 0 \Rightarrow (x+2)(x-2) = 0 \Rightarrow x = -2, 2$

Roots: $\alpha = -2, \beta = 2$

3. Region Test on $(x+2)(x-2) \leq 0$ **Test Box**



4. $\therefore -2 \leq x \leq 2$

QUESTION 2: If $(x-a)^2$ is a factor of $x^3 + 3px + q$ show that $p = -a^2$.

SOLUTION

You can prove this result by using the division process.

$$\begin{array}{r}
 x^2 - 2ax + a^2 \overline{) x^3 + 0x^2 + 3px + q} \\
 \underline{\mp x^3 \pm 2ax^2 \mp a^2x} \\
 2ax^2 + (3p - a^2)x + q \\
 \underline{\mp 2ax^2 \pm 4a^2x \mp 2a^3} \\
 (3p + 3a^2)x + (q - 2a^3)
 \end{array}$$

As $(x-a)^2$ is a factor, the remainder is zero, i.e. $0x + 0$.

$$\Rightarrow 3p + 3a^2 = 0 \Rightarrow p + a^2 = 0 \Rightarrow p = -a^2$$