

CALCULUS OPTION (Q 8, PAPER 2)

1999

8 (a) Use integration by parts to find $\int xe^x dx$.

(b) $f(x) = f(0) + \frac{f'(0)x}{1!} + \frac{f''(0)x^2}{2!} + \frac{f'''(0)x^3}{3!} + \dots$ is the Maclaurin series.

Derive the Maclaurin series for $f(x) = \cos x$ up to and including the term containing x^6 .

Hence write down the general term of the Maclaurin series for $f(x) = \cos x$.

Use the ratio test to show that the series converges for all $x \in \mathbf{R}$.

(c) A solid cylinder of radius r and height h has a fixed volume K .

(i) Express h in terms of r , π and K .

(ii) Find the ratio $r:h$ when the total surface area of the cylinder is a minimum.
Give your answer as a ratio of natural numbers.

SOLUTION

8 (a)

PARTS FORMULA $\int u dv = uv - \int v du$ **1**

STEPS

1. Call the original integral I (ignore limits of integration).
2. Let u equal the higher function in the list and find du by differentiation;
Let dv equal what is left and find v by integration.
NOTE: LIATE helps you to remember the order.
3. Substitute into Parts Formula. You will now be left with $\int v du$. You will either be able to integrate this integral normally or you must integrate by parts again.
4. If there are limits of integration, do them at the end.

LIST of Functions

1. **L**og
2. **I**nverse Trig
3. **A**lgebraic
4. **T**rigonometry
5. **E**xponential

1. $I = \int xe^x dx$

2.
$$\begin{array}{ll} u = x & dv = e^x dx \\ du = dx & v = e^x \end{array}$$

3. $I = x(e^x) - \int e^x dx$

$\therefore I = xe^x - e^x + c$

8 (b)

THE MACLAURIN FORMULA

$$f(x) = \frac{f(0)x^0}{0!} + \frac{f'(0)x^1}{1!} + \frac{f''(0)x^2}{2!} + \frac{f'''(0)x^3}{3!} \dots + \frac{f^{(n)}(0)x^n}{n!} + \dots \quad \text{3}$$

$$f(x) = \cos x \Rightarrow f(0) = 1$$

$$f'(x) = -\sin x \Rightarrow f'(0) = 0$$

$$f''(x) = -\cos x \Rightarrow f''(0) = -1$$

$$f'''(x) = \sin x \Rightarrow f'''(0) = 0$$

$$f^{(4)}(x) = \cos x \Rightarrow f^{(4)}(0) = 1$$

$$f^{(5)}(x) = -\sin x \Rightarrow f^{(5)}(0) = 0$$

$$f^{(6)}(x) = -\cos x \Rightarrow f^{(6)}(0) = -1$$

$$\cos x = \frac{1x^0}{0!} + \frac{0x^1}{1!} - \frac{1x^2}{2!} + \frac{0x^3}{3!} + \frac{1x^4}{4!} + \frac{0x^5}{5!} - \frac{1x^6}{6!}$$

$$\therefore \cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!}$$

STEPS TO FIND GENERAL TERM

1. The powers and coefficients of each series are in an arithmetic series. Use the formula for the general term of an arithmetic series T_n to generate u_n .

$$T_n = a + (n-1)d \quad \dots \quad \text{4}$$

2. Sometimes the signs alternate: +, -, +, -, +, -..... Multiply by $(-1)^{n-1}$ to achieve this alternation.

Powers, Factorial: 0, 2, 4,..... [Arithmetic series $a = 0, d = 2$]

$$T_n = 0 + (n-1)2 = 2n - 2$$

Signs alternate.

Therefore, general term for $\cos x$: $u_n = (-1)^{n-1} \frac{x^{2n-2}}{(2n-2)!}$

$$\sum_{n=1}^{\infty} u_n \text{ is } \mathbf{convergent} \text{ if } \lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| < 1. \text{ It is } \mathbf{divergent} \text{ if } \lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| > 1. \quad \dots \quad \text{2}$$

STEPS

1. Read off u_n from $\sum_{n=1}^{\infty} u_n$.
2. Find u_{n+1} .
3. Evaluate $\lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right|$. If $\lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| < 1$ the series is **convergent**. If

$\lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| > 1$ the series is **divergent**. If $\lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| = 1$ the test is **inconclusive**.

$$1. u_n = \frac{(-1)^{n-1} x^{2n-2}}{(2n-2)!}$$

$$2. u_{n+1} = \frac{(-1)^n x^{2n}}{(2n)!}$$

$$3. \lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(-1)^n x^{2n}}{(2n)!} \times \frac{(2n-2)!}{(-1)^{n-1} x^{2n-2}} \right| = \lim_{n \rightarrow \infty} \left| \frac{(-1)x^2}{(2n)(2n-1)} \right| = 0 < 1$$


Therefore, the series is convergent for all $x \in \mathbf{R}$.

8 (c)

STEPS

1. Identify the quantity to be maximised/minimised and give it a suitable symbol. **Example:** V for volume.
2. Draw a diagram (if necessary) and put in the variable(s).
3. Write the quantity in terms of this/these variable(s).
4. If there are 2 variables get rid of one in terms of the other using extra information.
5. Hence, write the quantity as a function of a single variable.
6. Differentiate the quantity with respect to the variable. Set it equal to zero and solve for the variable.
7. Substitute the value of the variable back into the quantity to find the maximum/minimum value.

1. Surface Area, A
2. Diagram shown.



$$3. A = 2\pi r^2 + 2\pi rh$$

$$4. K = \pi r^2 h \Rightarrow h = \frac{K}{\pi r^2}$$

$$5. A = 2\pi r^2 + 2\pi r \left(\frac{K}{\pi r^2} \right)$$

$$\Rightarrow A = 2\pi r^2 + 2 \left(\frac{K}{r} \right)$$

$$\therefore A = 2\pi r^2 + 2Kr^{-1}$$

$$6. \frac{dA}{dr} = 4\pi r - 2Kr^{-2} = 4\pi r - \frac{2K}{r^2}$$

$$\frac{dA}{dr} = 0 \Rightarrow 4\pi r = \frac{2K}{r^2}$$

$$\Rightarrow 2r = \frac{K}{\pi r^2}$$

$$\Rightarrow 2r = h$$

$$\Rightarrow \frac{r}{h} = \frac{1}{2}$$

$$\therefore r : h = 1 : 2$$

ANSWERS:

8 (c) (i) $\frac{K}{\pi r^2}$ (ii) 1:2